











## ACKNOWLEDGMENTS

We thank Giulia Di Vito (L'Aquila) for field and lab work. Thanks are also due to Paola Lombardo (Limnoconsulting, Rome) for English language revision and useful comments on a first draft of the manuscript. Comments and suggestions of the editor and two anonymous reviewers substantially improved the text. The research was partially funded by the Regione Abruzzo (DPC024 - Servizio Gestione e Qualità delle Acque).

## REFERENCES

- Anderson MJ, 2001. A new method for non-parametric multivariate analysis of variance. *Aust. Ecol.* 26:32-46.
- Barquín J, Death RG, 2006. Spatial patterns of macroinvertebrate diversity in New Zealand springbrooks and rithral streams. *J. N. Am. Benthol. Soc.* 25:768-786.
- Barquín J, Death RG, 2009. Physico-chemical differences in karst springs of Cantabria, northern Spain: Do invertebrate communities correspond? *Aquat. Ecol.* 43:445-455.
- Botosaneanu L, 1998. *Studies in crenobiology. The biology of springs and springbrooks.* Backhuys Publishers, Leiden: 261 pp.
- Bottazzi E, Bruno MC, Pieri V, Di Sabatino A, Silveri L, Carolli M, Rossetti G, 2011. Spatial and seasonal distribution of invertebrates in Northern Apennine rheocrene springs. *Limnol.* 70:77-92.
- Cantonati M, Bertuzzi E, Spitale D, 2008. *The spring habitat. Biota and sampling methods.* MUSE, Trento: 350 pp.
- Cantonati M, Füreder L, Gerecke R, Jüttner I, Gössler J, 2012. Crenic habitats, hotspots for freshwater biodiversity conservation: toward an understanding of their ecology. *Freshw. Sci.* 31:463-480.
- Carroll TM, Thorp JH, 2014. Ecotonal shifts in diversity and functional traits in zoobenthic communities of karst springs. *Hydrobiologia* 738:1-20.
- Di Sabatino A, Cicolani B, Gerecke R, 2003. Biodiversity and distribution of water mites (Acari, Hydrachnidia) in spring habitats. *Freshwater Biol.* 48:2163-2173.
- Di Sabatino A, Cristiano G, Pinna M, Lombardo P, Miccoli FP, Marini G, Vignini P, Cicolani B, 2014. Structure, functional organization and biological traits of macroinvertebrate assemblages from leaf-bags and benthic samples in a third-order stream of Central Apennines (Italy). *Ecol. Indic.* 46:84-91.
- Di Sabatino A, Cristiano G, Pinna M, Lombardo P, Miccoli FP, Marini G, Vignini P, Cicolani B, 2015. Corrigendum to: Structure, functional organization and biological traits of macroinvertebrate assemblages from leaf-bags and benthic samples in a third-order stream of Central Apennines (Italy) [*Ecol. Indic.* 2014;46:84-91]. *Ecol. Indic.* 49:260.
- Di Sabatino A, Cristiano G, Di Sanza D, Lombardo P, Giansante C, Caprioli R, Vignini P, Miccoli FP, Cicolani B, 2016. Leaf-Nets (LN): a new quantitative method for sampling macroinvertebrates in non-wadeable streams and rivers. *Rivers Res. Appl.* 32:1242-1251.
- Friberg N, Bonada N, Bradley DC, Dunbar MJ, Edwards FK, Grey J, Hayes RB, Hildrew AG, Lamouroux N, Trimmer M, Woodward G, 2011. *Biomonitoring of human impacts in freshwater ecosystems: the good, the bad and the ugly*, p. 1-68. In: G. Woodward (ed.), *Ecosystems in a human-modified landscape.* Elsevier, Amsterdam.
- Gerecke R, Cantonati M, Spitale D, Stur E, Wiedenbrug S, 2011. The challenges of long-term ecological research in springs in the northern and southern Alps: indicator groups, habitat diversity, and medium-term change. *J. Limnol.* 70:168-187.
- Gerecke R, Maiolini B, Cantonati M, 2007. Collecting meio- and macrozoobenthos in springs, p. 265-274. In: M. Cantonati, E. Bertuzzi, and D. Spitale (eds.), *The spring habitat: biota and sampling methods.* MUSE, Trento.
- Glazier DS, 1991. The fauna of North American temperate cold springs: patterns and hypotheses. *Freshwater Biol.* 26:527-542.
- Glazier DS, 2014. *Springs.* In: E.A. Elias (ed.), *Reference module in earth systems and environmental sciences.* Elsevier, Amsterdam.
- Gooch JL, Glazier DS, 1991. Temporal and spatial patterns in mid-Appalachian springs. *Mem. Ent. Soc. Can.* 155:29-49.
- Humphries P, Gross JE, Serafini LG, Hawking JH, Chick AJ, Lake PS, 1998. Macroinvertebrate sampling methods for lowland Australian rivers. *Hydrobiologia* 364:209-218.
- Merritt RW, Cummins KW, 1996. *An introduction to the aquatic insects of North America.* Kendall Hunt Publishing Co, Debuque.
- Pinna M, Marini G, Mancinelli G, Basset A, 2014. Influence of sampling effort on ecological descriptors and indicators in perturbed and unperturbed conditions: A study case using benthic macroinvertebrates in Mediterranean transitional waters. *Ecol. Indic.* 37:27-39.
- Rosati M, Cantonati M, Fenoglio S, Segadelli S, Levati G, Rossetti G, 2016. Is there an ideal protocol for sampling macroinvertebrates in springs? *J. Freshwater Ecol.* 31:199-209.
- Sangiorgio F, Glazier DS, Mancinelli G, Basset A, 2010. How can habitat size influence leaf litter decomposition in five mid-Appalachian springs (USA)? The importance of the structure of the detritivorous guild. *Hydrobiologia* 654:227-236.
- Smith H, Wood PJ, Gunn J, 2003. The influence of habitat structure and flow permanence on invertebrate communities in karst spring systems. *Hydrobiologia* 510:53-66.
- Staudacher K, Füreder L, 2007. Habitat complexity and invertebrates in selected Alpine springs (Schütt, Carinthia, Austria). *Internat. Rev. Hydrobiol.* 92:465-479.
- Tachet H, Bournaud M, Richoux P, Usseglio-Polatera P, 2000. [Invertébrés d'eau douce: Systématique, biologie, écologie]. [Book in French]. CNRS Editions, Paris: 608 pp.
- Tiegs SD, Peter FD, Robinson CT, Uehlinger U, Gessner MO, 2008. Leaf decomposition and invertebrate colonization responses to manipulated litter quantity in streams. *J. N. Am. Benthol. Soc.* 27:321-331.
- von Fumetti S, Nagel P, 2011. A first approach to a faunistic crenon typology based on functional feeding groups. *J. Limnol.* 70: 47-154.
- von Fumetti S, Nagel P, Baltes B, 2007. Where a springhead becomes a springbrook - a regional zonation of springs. *Fund. Appl. Limnol.* 169:37-48.
- Williams DD, Danks HV, 1991. Arthropods of springs, with particular reference to Canada. *Mem. Ent. Soc. Can.* 155:1-135.